VOL. 499 JOURNAL OF CHROMATOGRAPHY JANUARY 19, 1990

CONTENTS

HONOUR VOLUME ON THE OCCASION OF THE 60TH BIRTHDAY OF Cs. HORVÁTH

Preface
by L. R. Snyder 1

Determination of competitive adsorption isotherms for modeling large-scale separations in liquid chromatography
by J. M. Jacobson (Brisbane, CA, U.S.A.) and J. Frenz (South San Francisco, CA, U.S.A.) 5

Quantitative comparison between the experimental band profiles of binary mixtures in overloaded elution chromatography and their profiles predicted by the semi-ideal model
by A. M. Katti (Knoxville, TN, U.S.A.) and G. Guiochon (Knoxville and Oak Ridge, TN, U.S.A.) 21

Scaling-up procedure from the range of milligrams to grams for the purification of amino acid derivatives in displacement chromatography
by F. Cardinali, A. Ziggiotti and G. C. Viscomi (Rome, Italy) 37

Purification of monoclonal antibodies by complex-displacement chromatography on CM-cellulose
by A. R. Torres and E. A. Peterson (Logan, UT, U.S.A.) 47

Preparative-scale high-performance liquid chromatographic separation and purification of 3′-azido-3′-deoxythymidine-5′-phosphate
by J. G. Turcotte, P. E. Pivarnik, S. S. Shirali, H. K. Singh and R. K. Sehgal (Kingston, RI, U.S.A.), D. Macbride (Wakefield, RI, U.S.A.) and N.-I. Jang and P. R. Brown (Kingston, RI, U.S.A.) 55

Thermodynamic model for electrostatic-interaction chromatography of proteins
by I. Mazsaroff (Cambridge, MA, U.S.A.) and L. Vàrady, G. A. Mouchawar and F. E. Regnier (W. Lafayette, IN, U.S.A.) 63

Physicochemical studies of biologically active peptides by low-temperature reversed-phase high-performance liquid chromatography
by D. E. Henderson and J. A. Mello (Hartford, CT, U.S.A.) 79

Reversed-phase chromatographic behavior of proteins in different unfolded states
by S. Lin and B. L. Karger (Boston, MA, U.S.A.) 89

High-performance liquid chromatography of amino acids, peptides and proteins. XCV. Thermodynamic and kinetic investigations on rigid and soft affinity gels with varying particle and pore sizes: comparison of thermodynamic parameters and the adsorption behaviour of proteins evaluated from bath and frontal analysis experiments
by F. B. Anspach, A. Johnston, H.-J. Wirth, K. K. Unger and M. T. W. Hearn (Clayton, Australia) 103

Factors influencing the performance of peptide mapping by reversed-phase high-performance liquid chromatography
by M. W. Dong (Norwalk, CT, U.S.A.) and A. D. Tran (Raritan, NJ, U.S.A.) 125

Binary and ternary salt gradients in hydrophobic-interaction chromatography of proteins
by Z. El Rassi, L. F. De Ocampo and M. D. Bacolod (Stillwater, OK, U.S.A.) 141

High-performance adsorption chromatography of transfer ribonucleic acids and proteins on 2-μm spherical beads of hydroxyapatite. Influence of sodium chloride and magnesium ions on the resolution
by J. Lindeberg, T. Srichaiyo and S. Hjertén (Uppsala, Sweden) 153

Unexpected elution behaviour of peptides with various reversed-phase columns
by H. Engelhardt (Saarbrücken, F.R.G.), G. Appelt (Leipzig, G.D.R.) and E. Schweinheim (Saarbrücken, F.R.G.) 165

Hydrophilic-interaction chromatography for the separation of peptides, nucleic acids and other polar compounds
by A. J. Alpert (Baltimore, MD, U.S.A.) 177

Investigation of some α-amylases by high-performance hydrophobic-interaction chromatography
by L. Szepesy, L. Vida, M. Tóth and E. László (Budapest, Hungary) 197

Analysis of glycoprotein-derived oligosaccharides by high-pH anion-exchange chromatography
by L. J. Basa and M. W. Spellman (South San Francisco, CA, U.S.A.) 205

Automated evaluation of tryptic digest from recombinant human growth hormone using ultraviolet spectra and numeric peak information
by H.-J. P. Sievert (Avondale, PA, U.S.A.) and S.-L. Wu, R. Chloupek and W. S. Hancock (South San Francisco, CA, U.S.A.) 221

Protein chromatography with pyridine- and alkyl-thioether-based agarose adsorbents
by S. Oscarsson and J. Porath (Uppsala, Sweden) 235

New latex-bonded pellicular anion exchangers with multi-phase selectivity for high-performance chromatographic separations
by J. R. Stillian and C. A. Pohl (Sunnyvale, CA, U.S.A.) 249

Micropellicular stationary phases for rapid protein analysis by high-performance liquid chromatography
by K. Kalghatgi (New Haven, CT, U.S.A.) 267

Comparison of retention behavior on polymeric resins and an alkyl-bonded silica phase in reversed-phase high-performance liquid chromatography
by S. Pedigo and L. D. Bowers (Minneapolis, MN, U.S.A.) 279

Physical and chemical characterization of microporous zirconia
by M. P. Rigney, E. F. Funkenbusch and P. W. Carr (Minneapolis, MN, U.S.A.) 291

Effect of coverage density on the retention mechanism in reversed-phase high-performance liquid chromatography
by B. Buszewski, Z. Suprynowicz and P. Staszczuk (Lublin, Poland) and K. Albert, B. Pfleiderer and E. Bayer (Tübingen, F.R.G.) 305

Studies on the stability of *n*-alkyl-bonded silica gels under basic pH conditions
by N. T. Miller and J. M. DiBussolo (Conshohocken, PA, U.S.A.) 317

Mechanism of retention in high-performance liquid chromatography on porous graphitic carbon as revealed by principal component analysis of structural descriptors of solutes
by R. Kaliszan and K. Ośmiałowski (Gdansk, Poland) and B. J. Bassler and R. A. Hartwick (Piscataway, NJ, U.S.A.) 333

Preparation of silicone-coated 5–25-μm I.D. fused-silica capillary columns for open-tubular liquid chromatography
by O. van Berkel-Geldof, J. C. Kraak and H. Poppe (Amsterdam, The Netherlands) . . . 345

Prediction of the high-performance liquid chromatographic retention behaviour of some benzodiazepine derivatives by thin-layer chromatography
by K. Valko, S. Olajos and T. Cserháti (Budapest, Hungary) 361

Study of partition models in reversed-phase liquid chromatography based on measured mobile phase solute activity coefficients
by W. J. Cheong and P. W. Carr (Minneapolis, MN, U.S.A.) 373

Retention characteristics of several compound classes in reversed-phase liquid chromatography with β-cyclodextrin as a mobile phase modifier
by R. M. Mohseni and R. J. Hurtubise (Laramie, WY, U.S.A.) 395

Reversed-phase ion-pair liquid chromatography of glucuronides. Retention and selectivity
by M. Stefansson and D. Westerlund (Uppsala, Sweden) 411

Basis of the rational selection of the hydrophobicity and concentration of the ion-pairing reagent in reversed-phase ion-pair high-performance liquid chromatography
by A. Bartha and G. Vigh (College Station, TX, U.S.A.) and Z. Varga-Puchony (Veszprem, Hungary) 423

Estimation of the reversed-phase liquid chromatographic lipophilicity parameter log k'_w using ET-30 solvatochromism
by J. J. Michels and J. G. Dorsey (Gainesville, FL, U.S.A.) 435

Strategies of mobile phase transfer from thin-layer to medium-pressure liquid chromatography with silica as the stationary phase
by S. Nyiredy, K. Dallenbach-Toelke, G. C. Zogg and O. Sticher (Zürich, Switzerland) . 453

Investigation and improvement of the Horváth–Lin equation of state
by P. Benedek and T. Jankó (Budapest, Hungary) 463

Reversed-phase liquid chromatographic studies of non-ionic surfactants and related solutes
by P. Varughese, M. E. Gangoda and R. K. Gilpin (Kent, OH, U.S.A.) 469

Effect of alcohol chain length, concentration and polarity on separations in high-performance liquid chromatography using bonded cyclodextrin columns
by I. Z. Atamna, G. M. Muschik and H. J. Issaq (Frederick, MD, U.S.A.) 477

Optimization of selectivity, detectability and analysis time in high-performance liquid chromatography
by S. Ahuja (Suffern, NY, U.S.A.) 489

Strategy for peak tracking in liquid chromatography on the basis of a multivariate analysis of spectral data
by J. K. Strasters, H. A. H. Billiet and L. de Galan (Delft, The Netherlands) and B. G. M. Vandeginste (Nijmegen, The Netherlands) 499

Peak tracking and subsequent choice of optimization parameters for the separation of a mixture of local anaesthetics by high-performance liquid chromatography
by J. K. Strasters, F. Coolsaet, A. Bartha, H. A. H. Billiet and L. de Galan (Delft, The Netherlands) 523

Performance of micro-liquid chromatographic columns in an industrial environment: a case history
by N. G. F. M. Lammers and J. H. M. van den Berg (Weesp, The Netherlands) and M. Verzele and C. Dewaele (Ghent, Belgium) 541

Direct coupling of capillary liquid chromatography with conventional high-performance liquid chromatography
by T. Takeuchi, M. Asai, H. Haraguchi and D. Ishii (Nagoya, Japan) 549

Appraisal of four pre-column derivatization methods for the high-performance liquid chromatographic determination of free amino acids in biological materials
by P. Fürst, L. Pollack, T. A. Graser, H. Godel and P. Stehle (Stuttgart, F.R.G.) 557

Preparation of electrophoric derivatives of N7-(2-hydroxyethyl)guanine, an ethylene oxide DNA adduct
by K. Allam, M. Saha and R. W. Giese (Boston, MA, U.S.A.) 571

Application of 3-(2-furoyl)quinoline-2-carbaldehyde as a fluorogenic reagent for the analysis of primary amines by liquid chromatography with laser-induced fluorescence detection
by S. C. Beale, Y.-Z. Hsieh, D. Wiesler and M. Novotny (Bloomington, IN, U.S.A.) . . . 579

Chromatographic studies on the binding, action and metabolism of (−)-deprenyl
by H. Kalász, L. Kerecsen, J. Knoll and J. Pucsok (Budapest, Hungary) 589

Determination of fluoxetine and norfluoxetine by high-performance liquid chromatography
by S. H. Y. Wong, S. S. Dellafera, R. Fernandes and H. Kranzler (Farmington, CT, U.S.A.) 601

On-line high-performance liquid chromatography for the determination of cephalosporin C and by-products in complex fermentation broths
by K. Holzhauer-Rieger, W. Zhou and K. Schügerl (Hannover, F.R.G.) 609

Determination of D-*myo*-1,2,6-inositol trisphosphate by ion-pair reversed-phase liquid chromatography with post-column ligand exchange and fluorescence detection
by H. Irth, M. Lamoree, G. J. de Jong, U. A. Th. Brinkman and R. W. Frei (Amsterdam, The Netherlands) and R. A. Kornfeldt and L. Persson (Perstorp, Sweden) 617

Separation of carotenes on cyclodextrin-bonded phases
by A. M. Stalcup, H. L. Jin and D. W. Armstrong (Rolla, MO, U.S.A.) and P. Mazur, F. Derguini and K. Nakanishi (New York, NY, U.S.A.) 627

Comparative supercritical-fluid chromatographic properties of carbon dioxide and sulphur hexafluoride–ammonia mixtures with packed columns
by J. L. Veuthey (Genève, Switzerland) and J. L. Janicot, M. Caude and R. Rosset (Paris, France) 637

Use of high-performance liquid chromatography for the characterization of synthetic copolymers
by G. Glöckner (Dresden, G.D.R.) and H. G. Barth (Wilmington, DE, U.S.A.) 645

Effect of operating parameters on polymer molecular weight accuracy with time-delay, exponential-decay thermal field flow fractionation
by J. J. Kirkland and W. W. Yau (Wilmington, DE, U.S.A.) 655

Ion chromatography of transition metals on an iminodiacetic acid bonded stationary phase
by G. Bonn and S. Reiffenstuhl (Innsbruck, Austria) and P. Jandik (Milford, MA, U.S.A.) 669

Calculation of column performance in nitrate removal from water supplies by anion exchange
by R.-J. Leu, Y.-L. Hwang and F. G. Helfferich (University Park, PA, U.S.A.) 677

Determination of inorganic anions by flow injection analysis and high-performance liquid chromatography combined with photolytic-electrochemical detection
by L. Dou and I. S. Krull (Boston, MA, U.S.A.) 685

Kinetics of cysteine oxidation in immobilized pH gradient gels
by M. Chiari, C. Chiesa, P. G. Righetti and M. Corti (Milan, Italy) and T. Jain and R. Shorr (Malvern, PA, U.S.A.) 699

Capillary electrophoresis with indirect amperometric detection
by T. M. Olefirowicz and A. G. Ewing (University Park, PA, U.S.A.) 713

High-efficiency liquid extractor for isolation of a desired material from complex organic mixtures
by M. Brenner, C. Moirandat and R. Schlimme (Basle, Switzerland) 721

Gas chromatographic study of the solution thermodynamics of organic solutes in tetraalkylammonium alkanesulfonate and perfluoroalkanesulfonate solvents
by R. M. Pomaville and C. F. Poole (Detroit, MI, U.S.A.) 749

Author Index 761

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*)